

Introduction: Localized Structures in Dissipative Media: From Optics to Plant Ecology

M. Tlidi¹, K. Staliunas², K. Panajotov^{3,4}, A.G. Vladimirov⁵ and M.G. Clerc^{6,1}

¹¹ *Département de Physique, Université Libre de Bruxelles (U.L.B.),
CP 231, Campus Plaine, B-1050 Bruxelles, Belgium.*

² *Departament de Física i Enginyeria Nuclear, Universitat Politècnica
de Catalunya, Colom 11, 08222 Terrassa (Barcelona), Spain.*

³ *Brussels Photonics Team, Department of Applied Physics and Photonics (B-PHOT TONA),
Vrije Universiteit Brussels, Pleinlaan 2, B-1050 Brussels, Belgium.*

⁴ *Institute of Solid State Physics, 72 Tzarigradsko Chaussee Blvd., 1784 Sofia, Bulgaria*

⁵ *Weierstrass Institute, Mohrenstrasse 39, D-10117 Berlin, Germany.*

⁶ *Departamento de Física, FCFM, Universidad de Chile, Blanco Encalada 2008, Santiago, Chile.*

Localised structures in dissipative appears in various fields of natural science such as biology, chemistry, plant ecology, optics and laser physics. The proposed theme issue is to gather specialists from various fields of non-linear science toward a cross-fertilisation among active areas of research. This is a cross-disciplinary area of research dominated by the nonlinear optics due to potential applications for all-optical control of light, optical storage, and information processing. This theme issue contains contributions from 18 active groups involved in localized structures field and have all made significant contributions in recent years.

PACS numbers:

I. INTRODUCTION

Localized structures are "dissipative structures" found far from equilibrium. The term "dissipative structure" has been introduced by Ilya Prigogine to describe the spontaneous appearance of periodic structures as a result of irreversible processes maintained by exchanges of matter and/or energy with a non-equilibrium environment [1, 2]. Dissipative structures generally evolve on macroscopic scales and can only be maintained by continuous application of a non-equilibrium constraint. The spontaneous emergence of dissipative structures arise from a principle of self-organization that can be either in space and/or in time. A classic example of spatial self-organization is provided in the context of chemical reaction diffusion system referred to as Turing instability [3]. This instability is characterized by a symmetry breaking that leads spontaneously to the formation of periodic structures with an intrinsic wavelength. In this case the wavelength is determined solely by the dynamical parameters such as diffusion coefficient and the inverse characteristic time associated with chemical kinetics. The emergence of dissipative structures with an extrinsic wavelength has been reported in an early report by Bénard [4]. Transition from conductive to convective regimes occurs in a plane horizontal layer of fluid heated from below, in which the fluid develops a regular pattern of convection cells referred to as Bénard cells. In this case, the wavelength, i.e., the size of the pattern is determined by the thickness of the fluid layer and not by the dynamical parameters. The first experimental evidence of dissipative structures with an intrinsic wavelength has been given by Castets et al.[5] and by Ouyang and Swin-

ney [6] by using CIMA (Chlorite-Iodide-Malonic-Acid) reaction. Earlier, the Soviet scientist Belousov has predicted the oscillating behavior and the ability of chemical waves concentration to spread [7, 8]. However, dissipative structures have not been discussed.

Spontaneous symmetry breaking and self-organisation phenomena appear in various field of nonlinear science such as nonlinear optics, fiber optics, plant ecology, fluid mechanics, granular matter, plant ecology. The link between the classical Turing instability and transverse patterns in nonlinear optics has been first established in the seminal paper by Lugiato and Lefever [9]. Dissipative structures could be localized in space and/or in time. Many driven systems exhibit localized structures (LSs), often called localized spots and localized patterns, which may be either isolated, randomly distributed, or self-organized in clusters forming a well-defined spatial pattern. Localized structures (LSs) are homoclinic solutions (solitary or stationary pulses) of partial differential equations. The conditions under which LSs and periodic patterns appear are closely related. Typically, when the Turing instability becomes sub-critical, there exists a pinning domain where localized structures are stable. This is a universal phenomenon and a well documented issue in various fields of nonlinear science, such as chemistry, plant ecology, and optics. The experimental observation of LSs in driven nonlinear optical cavities has motivated further the interest in this field of research. In particular, LSs could be used as bits for information storage and processing. Several overviews have published on this active area of research [10–30]

Collectively, the papers in this Special Issue provide an overview of recent advances in the formation of Localized structures in dissipative media. This special issue

includes a selection of papers presented at the conference of "Dynamics days", Madrid 2013. It concerns various topics of localized structures in dissipative media which are briefly discussed below.

II. LOCALIZED STRUCTURES AND PATTERN FORMATION IN NONLINEAR OPTICS

Before the invention of lasers, the optics was linear (with some small exceptions, like Raman scattering), and no nonlinear dynamical regimes were ever considered in optics. As the spontaneous pattern formation can occur in strongly nonlinear systems driven far away from equilibrium, no pattern formation was expected in optics. First ideas on the spatial pattern formation in optics appeared in early 70-ties, decade after the laser invention, when the relation between coherent laser field in laser resonators and fluids/superfluids was recognized, based on the derivation of complex Ginzburg-Landau equation for the light fields [31]. Also filamentation of strong light fields in self-focusing media (in analogy to singularity appearance in plasmas) [32] can be considered as the very initial works on the nonlinear light pattern formation. In spite of these initial ideas the real pattern forming in optics was considered unrealistic, mainly because of the weakness of nonlinearities there. The real interest in nonlinear optical patterns started in the late 80-ies and early 90-ies. In [9, 33, 34], a nontrivial nonlinear transverse mode formation in laser beams were shown. In [35], it was recognized that the nonlinear laser equations (Maxwell-Bloch equations) admit vortex solutions, which were soon confirmed experimentally [36]. These pioneering works were followed by an increasing number of investigations, mostly attempting to understand the mechanisms of nonlinear pattern formation by deriving reduced models for a laser and other broad aperture nonlinear resonators, such as modified Ginzburg-Landau equation [37], and Swift-Hohenberg equation [38–42]. Such approach has been applied for other types of nonlinear resonators filled with photorefractive crystal [43], with parametric nonlinearity [44, 45], also passive driven nonlinear-Kerr media [12, 46]. The number of works on different aspects of transverse pattern formation was exploding in 90-ies (see [10–14]). Very different issues have been considered, which are impossible to review in this short introduction. To mention a few, three-dimensional patterns in resonators (localized light bullets) [47–54], patterns in "rocked" resonators [55–57], and hyperbolic patterns [58, 59]. Exotic phenomena, like optical vortices and vortex ensembles [35, 36, 60–64], which can show chaotic dynamics (vortex mediated turbulence) [64, 65], and vortices that arrange themselves in regular lattices [66–70] have been demonstrated. Nonlinear patterns also appear in other optical configurations such as: mirrorless configuration with unidirectional light propagation in nonlinear

media, leading to filamentation effects, to formation of bright solitons [71], and optical vortices [72, 73]; counterpropagating configuration leading to hexagons [74]; systems with feedback loop, showing a large variety of extended patterns [75, 76]. Localized structures were reported in bistable nonlinear resonators such as lasers with saturable absorbers) [11, 77, 78], in passive nonlinear resonators [79, 80], and optical parametric oscillators [81, 82]. Phase solitons have been demonstrated far from any pattern forming instability [83–86].

III. LOCALIZED STRUCTURES IN VCSELS

Owing to their large Fresnel number and short cavity, Vertical-Cavity Surface-Emitting Lasers (VCSELs) are best suited for cavity soliton (CS) studies and potential applications (for recent reviews see [25, 87]). VCSELs were first suggested and realized in 1979 [88] and it took more than 10 years to bring them to comparable performance to edge-emitting lasers [89, 90]. Nowadays VCSELs are replacing edge-emitting lasers in short and medium distance optical communication links thanks to their inherent advantages: much smaller dimensions, circular beam shape that facilitates coupling to optical fibers, two-dimensional array integration and on wafer testing that brings down the production cost [90]. As VCSELs emit light perpendicular to the surface and the active quantum wells, their cavity length is of the order of $1\text{ }\mu\text{m}$ - the wavelength of the generated light. Thanks to the maturity of the semiconductor technology VCSELs can be made homogeneous over a size of hundreds of μms while the characteristic CS size is about $10\text{ }\mu\text{m}$. Furthermore, the timescales of the semiconductor laser dynamics and CS formation are in the ns scale, which allows for fast and accurate gathering of data. Finally, VCSEL physics and dynamics are quite well understood [90–92]. Therefore, many theoretical and experimental studies on CS formation in VCSELs have been carried out [93]–[106]. CSs have been successfully demonstrated in broad area VCSELs both below [93, 95] and above [96] the lasing threshold when injecting a holding beam of appropriate frequency. Spatially localized structure has also been found out in medium size VCSELs but only by using their particular polarization properties [97]. CS lasers (CSLs) in a VCSEL system without a holding beam have been demonstrated both experimentally [98] and theoretically [99] in VCSELs subject to frequency selective optical feedback and in face to face coupled VCSELs [100, 101]. In these systems, the VCSELs are placed in self imaging optical systems with either an external grating or another VCSEL biased below lasing threshold, so that the system becomes bistable. Lasing spots spontaneously appear in these systems and can be switched on and off by another laser beam. As a matter of fact broad area laser with saturable absorber has been the first system in which CS

have been predicted and studied theoretically [102–104]. CSs in a monolithic optically pumped VCSEL with a saturable absorber have been demonstrated in [105] and their switching dynamics studied in [106]. Several applications of CSs in VCSELs have been demonstrated: optical memory [107], optical delay line [108] and optical microscopy [109].

IV. LOCALIZED STRUCTURES IN PHOTONIC CRYSTALS, LEFT-HANDED MATERIALS AND EXCITONS-POLARITONS

High-speed switching i.e., processing speed in optical systems combined with possibility of parallel processing in broad area systems gives a clear advantage of the optical systems with respect to electronic circuits. However, the optical systems have also a serious disadvantage with respect to electronic systems - the relatively large spatial extent of several wavelengths, i.e. of tens of micrometers. This is a serious disadvantage limiting application of nonlinear optical patterns in microprocessing technologies. Therefore, the straightforward application of cavity solitons for optical memory and all-optical processing has never come to practice. Nevertheless, the advance of micro- and nano- fabrication technologies offers new possibilities for optical pattern: novel micro-modulated materials, like photonic crystals or nanomaterials that allow the manipulation of light propagation (zero or negative refraction and diffraction), thus suppressing the diffractive broadening of the beams, and strongly reducing the size of the nonlinear photonic structures [110–112]. On the other hand, the combined light-matter structures, the so called excitons and surface polaritons, are also of drastically reduced dimensions, due to light-matter coupling [113–115].

The fundamental limitations of LSs is their spatial size, which is due to the diffraction phenomenon in optical resonators. To overcome this limit, use of left-handed metamaterials is proposed, i.e. engineered materials with simultaneously negative permittivity and permeability [116–120]. These materials were first used by the scientific community to realize imaging systems with sub-wavelength resolution then to target potential applications including invisibility cloaks and perfect optical concentrators. They also allow for phase compensation in optical structures containing both left- and right-handed materials. Recently, it has been shown that such materials could be used to improve telecommunication systems by lowering power threshold and increasing data density in optical memories [121–124]. More recently, an experimental evidence of localized structure in left-handed materials has been provided [125].

V. FRONT DYNAMICS IN SPATIALLY EXTENDED SYSTEMS

Macroscopic systems under the influence of injection and dissipation of quantities such as energy and momenta usually exhibit coexistence of different states, which is termed multistability. This is clearly a demonstration that macroscopic systems are of nonlinear nature. Heterogeneous initial conditions usually caused by the inherent fluctuations generate spatial domains, which are separated by their respective interfaces. These interfaces are known as front solutions or interfaces or domain walls, etc. [14]. Interfaces between these metastable states appear in the form of propagating fronts and give rise to a rich spatiotemporal dynamics [126–128]. Front dynamics occurs in systems as different as walls separated magnetic domains [129], directed solidification process [130], nonlinear optical systems [131–134], oscillating chemical reactions [135], fluidized granular media [136–141, 141], or population dynamics [18, 142, 143]. From the point of view of dynamical system theory, in one spatial dimension a front is a nonlinear solution that is identified in the comoving frame system as a heteroclinic orbit linking two spatially extended states [144, 145]. The dynamics of the interface depends on the nature of the states that are connected. In the case of a front connecting a stable and an unstable state, it is called a Fisher-Kolmogorov-Petrovsky-Piskunov (FKPP) front [18, 146, 147]. One of the characteristic features of these fronts is that the speed they move is not unique, nonetheless determined by the initial conditions. When the initial condition is bounded, after a transient, two counter propagative fronts with the minimum asymptotic speed emerge [18, 147]. In case that the nonlinearities are weak, this minimum speed is determined by the linear or marginal-stability criterion and fronts are usually referred to as pulled [147]. In the opposite case, the asymptotic speed can only be determined by nonlinear methods and fronts are referred to as pushed [147]. Variational methods are particularly efficient to determine appropriate approaches for the minimum speed for pushed fronts [148]. FKPP fronts have been observed in Taylor-Couette [149], Rayleigh-Benard experiments [150], pearling and pinching on the propagating Rayleigh instability [151], spinodal decomposition in polymer mixtures [152] and liquid crystal light valve [153], to mention a few.

The above scenario changes completely for a front connecting two stable states. In the case of two uniform states, a gradient system tends to develop the most stable state, in order to minimize its energy, so that the front always propagates toward the most energetically favorable state. It exists only as one point in parameter space for which the front is motionless, which is usually called the Maxwell point, and is the point for which the two states have exactly the same energy [154]. The evolution of front solutions can be regarded as a particle-type

one, i.e., they can be characterized by a set of continuous parameters such as position, width of core and so forth. Close to the Maxwell point, based on the method of variation of parameters, one can analytically determine the speed of the front [126]. For variational systems away from the Maxwell point one can have implicit expressions for the front speed [126]. In non-variational case, the analytical expression of the front speed is a problem still unresolved.

System with discrete reflection symmetry possess two equivalent states with interfaces, or domain walls which are generically at rest. Indeed, the two connected states are "energetically" equivalent. These front solutions are termed kinks. However, under spontaneous breaking of the parity symmetry, these fronts can acquire a nonzero asymptotic speed. This phenomenon is the denominated Ising-Bloch transition [155] and has been observed in systems as different as ferromagnetic systems [156, 157], liquid crystals [158, 159], chemical reactions [160], and nonlinear optical cavities [161]. Gradient or variational systems do not exhibit this phenomenon, because the front speed is proportional to the energy difference between the two equivalent states. A different situation is that of a front connecting a homogeneous and a periodic state. Indeed, the existence of a pinning range for a front between these states was predicted by Pomeau [126]. In this case, a pinning-depinning phenomenon is expected to occur as a result of the competition between a symmetry breaking of the global energy that tends to favor the front propagation in one direction and spatial modulations that tend to block the front by introducing local potential barriers at the front core dynamics [162, 163]. Depending on the dominating effect, the front can either stay motionless- blocked over a large range, therefore called the pinning range-or propagate with periodic leaps apart from it. Starting from a critical value of the control parameter, the pinning-depinning transition occurs by a loss of stability of the pinned front.

The existence of a pinning range, and associated pinning-depinning phenomenon, has a fundamental relevance in numerous domains where front propagation is involved. As examples, we can cite vibrated fluids [164], chemical reactions [135, 165], microfluidic chips [166], wetting of micro-structured surfaces [167], control of the motility of bacteria [168], and growth of self-assembly monolayers [169]. Indeed, spatial discreteness and spatial modulations can be seen as the bases for a wealth of life behaviors, where the emergence of complexity results from the microscopic granularity of the system [170]. As for the experimental studies of the pinning-depinning phenomenon, only a few approaches have been proposed up to date. In a two-dimensional spatially forced system, an experimental characterization of front propagation has shown the anisotropy of the front velocity [171]; however, the issue of a pinning range was not addressed. Only recently, by employing a 1D spatially forced liquid-

crystal system, the experimental demonstration of the existence of a pinning range has been achieved [172].

VI. TEMPORAL LOCALIZED STRUCTURES

In the temporal domain, localized structures have also been observed in passive fiber resonators. These devices constitute a basic configuration in nonlinear fiber optics. More specifically, experimental studies have demonstrated that when these resonators are pumped by a continuous wave, they exhibit spontaneously self-organized temporal structures in the form of trains of short pulses with well defined repetition rate [173–175]. The theoretical description of all fiber resonators is provided by the Lugiato and Lefever model (LL model, [176]). The breakup of continuous wave into trains of pulses is attributed to the competition between the nonlinear mechanism which is originated from the intensity-dependent refractive index (Kerr effect) that tends to amplify locally the field intensity, the chromatic dispersion which on the contrary tends to restore uniformity, and dissipation.

Besides a pulse train distribution, temporal localized structures (TLS) are found in a well-defined region of parameters called a pinning zone. In this regime, the system exhibits a coexistence between two states: the uniform background and the train of pulses of light that emerges from subcritical modulational instability [80]. They have been experimentally observed in fiber ring resonator [177]. Theoretical predictions and experimental observations of temporal localized structures using standard optical fibers allow for the conception of all-optic systems for generation of signals with high repetition rate or to operate as all-optical memories with capacity that can reach 25 Gbits/s [177]. In addition to temporal localised structures, front dynamics have been analytically predicted and experimentally observed in a simple passive fiber cavity synchronously pumped by a pulsed laser [178]. Synchronous pumping means that the time-of-flight of the light pulses in the cavity is adjusted to the laser repetition time. Recently, an asymptotic behavior of the front velocity has been established near the up-switching point of the bistable response curve [179]. The front dynamics obeys a generic power law when the front velocity approaches asymptotically its linear growing value as predicted by van Saarloos [147].

When all fiber cavity is operating close to the zero dispersion wavelength, high-order chromatic dispersion effects may play an important role in the dynamics of a photonic crystal fiber (PCF) [180–182]. PCFs allow for a high control of the dispersion curve and permit exploring previously inaccessible parameter regimes [183, 184]. The inclusion of the fourth order dispersion allows the modulational instability to have a finite domain of existence delimited by two pump power values [185] for the stabilization of dark temporal cavity solitons in PCF res-

onators [186]. Together with this effect, the third order dispersion causes a spontaneous broken reflection symmetry and allows the motion of both periodic and localized structures that are generated in the resonator [187].

VII. VEGETATION PATTERN IN SEMI(ARID) ECOSYSTEMS

Desertification is one of the grand environmental challenges in EU and in the world. In last decade, the concepts of symmetry breaking and self-organization have been applied to plant ecology systems where dissipative structures are generically called vegetation patterns. A well-documented example is "tiger bush". The observation of this phenomenon was possible thanks to the development of aerial photography in the early forties. Vegetation patterns were described for the first time 64 years ago by Macfadyen in British Somaliland [188]. In this paper, Macfadyen states that:

The study of the first two patterns, of which I have been able to find no previous recognition, is not conveniently pigeon-holed under any one accepted branch of knowledge, and the phenomena are thus awkward to classify. They are manifestly within the province of botany and ecology; the essential background concerns geomorphology and meteorology; the causes as I believe, must be investigated by physics and mathematics. [188]

Since then, vegetation patterns have been observed and studied in many regions throughout the world [189]. Vegetation patterns are sparsely populated or bare areas alternating with dense vegetation patches. They are made of either grass or trees and shrubs. By increasing the aridity, the spatial distribution (periodic or aperiodic) of patches first displays bare spots, then stripes, and finally spots. The spatial distribution occurs on a decametric to hectometric scale. This large scale phytosociological self-organization leading to nonuniform and non random distribution of the phytomass is the rule rather than the exception. They were observed on vast territories of many arid and semiarid regions of Africa, Middle East, Australia and North America. The terms arid and semiarid refer to climatic conditions where the water and nutrient resources are scarce; typically, in regions where the annual rainfall (50 – 750 mm) is low compared to the potential of evapo-transpiration ($PET; 1.5 \times 10^3$ mm).

Mathematical modeling of the vegetation growth constitutes an important tool towards the understanding of the mechanism leading to the desertification phenomenon in semiarid and arid landscapes. To explain the origin of the vegetation patterns three main mechanisms have been proposed. The following classification of ecological models was suggested by Lefever and Turner in [190]:

(i) The non-local FKPP (Fisher, Kolmogorov, Petrovsky, Piskunov) approach: it focuses on the relationship between the structure of individual plants and the

facilitation-competition interactions existing within plant communities. The biomass density is defined at the individual plant level; the modeling calls for no other state variable [9, 191–193].

(ii) The reaction-diffusion approach: it emphasizes the influence on vegetation patches of water transport by below ground diffusion and/or above ground run-off. The biomass density is defined at the patch level. Together with the water concentration below ground and/or in some surface ground layer, it constitutes the set of state variables [194–199].

(iii) The stochastic approach: it focuses on the "constructive" role of environmental randomness as a source of noise induced symmetry breaking transitions triggering pattern formation. These transitions occur when the variance of environmental noise increases while its average value remains constant [28, 200].

Besides the behavior of periodic vegetation patterns, the same self-organization mechanism predicts the existence of aperiodic, localized structures. They consist of localized patches of vegetation, randomly distributed on bare soil [201–203] or, on the contrary, consist of localized spots of bare soil, randomly distributed in an otherwise uniform vegetation cover [193]. Well documented examples are the fairy circles existing in the Namib desert. These circular areas devoid of any vegetation can reach diameters of up to 14 m, which exceeds by more than one order of magnitude the size of the tall grasses surrounding them. In a recent study, van Rooyen *et al.* [204] have investigated in depth the strengths and shortcomings of several hypotheses concerning their origin. In this respect, the authors have been able to rule out external causes such as a possible existence of localized radioactive areas unsuited for the development of plants or a link between fairy circles and the activity of termites or the proximity of their nests; among the other hypotheses considered by the authors (notably, the release of allelopathic compounds and the possible existence of stimulatory or inhibitory influences due to interactions between different plant species), none was found which fitted satisfactorily their experimental observations. Recent progress has been made in the modeling of fairy circles [193, 205–209]. The origin of this phenomenon is still matter of debate.

VIII. CONTRIBUTIONS TO THE THEME ISSUE

The contributions to this theme issue are devoted to the investigation of the formation, stability and control of localized structures in various research areas: nonlinear optics (including laser dynamics and fiber optics), fluid mechanics, chemistry and plant ecology. The results presented in the contributions indicate that considerable progress has been made in understanding of the

properties of non-linear localised structures in dissipative systems. Most of contributions deal with non-linear photonics devices having potential applications for all-optical control of light, optical storage, and information processing. The contributions are grouped by research area. The paper by Vladimirov et al. [210] presents the first experimental evidence of localized structures of light in a medium size bottom-emitting InGaAs multiple quantum well VCSEL operated in injection-locked regime. A scheme based on delay feedback to control the self-mobility of such localized structures was proposed and studied theoretically. In the case of two coupled VCSELs, Turconi et al. [211] perform a statistical analysis of time-resolved commutation experiments and investigate the role of wavelength, polarization and pulse energy in the switching process. Vahed et al. [212] investigate periodic and chaotic localized structures in VCSEL with a saturable absorber. Using a nematic liquid crystal cells with a photosensitive wall and negative dielectric anisotropy, Barboza et al [213] demonstrate experimentally and numerically the possibility of generation of stable vortices with swirling arms. The demonstration and the characterization of families of moving solitons in lattices composed of metallic nanoparticles with nonlinear Kerr-like response and an external driving field is provided by Noskov et al. [214]. A system formed by an array of plasmonic waveguides, with gain applied to selected hot spot cores is discussed by Ding et al. [215]. A review on localized modes in dissipative lattice media is presented by Malomed et al. [216]. Another review on the dynamics of particles, waves as well as spatial and temporal localized structures in cavities formed by oscillating mirrors is contributed by Rosanov [217]. Various collisions of 2D-localized structures are investigated and a possibility of breather formation as a result of collision is presented. The contribution by de Varcарcel et al. [218] presents a review on rocking phenomenon in various spatially extended systems and discuss their applications in optics. A Nagumo-Kuramoto model exhibiting a localised spatiotemporal chaotic type of behavior often called chaoticon is discussed by Verschueren et al. [219]. The mechanism responsible for the appearance of such chaotic regime is attributed to front interactions. Tesio et al. [220] discuss the self-organisation phenomenon in cold atomic gases that manifests itself as a spontaneous emergence of long-range spatial order from fluctuations in the transverse plane. Using the Kuramoto model, this process is interpreted as a synchronisation transition in a fully connected network of fictitious oscillators. In non-linear fiber optics, a vector model describing an erbium doped mode-locked fibre laser with carbon nanotube saturable absorber is investigated by Sergeyev [221]. In particular, the author discusses the formation of a new family of vector dissipative solitons with fast and slowly evolving polarization states in this system. Bahloul et al. [222] perform analytical and numerical studies of the

combined influence of third and fourth order dispersion on the dynamics of temporal localized structures in photonic crystal fiber cavity. This analysis reveals the existence of moving temporal localized structures of light.

Chabchoub et al. [223] provide for the first time evidence of localised structures in the form of Kuznetsov-Ma soliton and Akhmediev breather in a water wave flume. Their analytical study based on nonlinear Schrödinger equation agrees with the experimental observations. The control of breathing localized structures by time-delayed optical feedback in a three-component reaction-diffusion system is investigated theoretically by Gurevich [224]. It is shown that varying the delay parameters can lead either to stabilization of the breathing or to delay-induced periodic and quasiperiodic oscillations of localized structures. Finally, three contributions on localized patches, fairy circles, and dissipative structures in plant ecology are presented in the end of the issue. They can be summarized as follows. The first contribution by Couteron et al. [225] presents plant morphologies observed in the Andean altiplano and interpret them in terms of localized structures leading to a patchy, aperiodic distribution of the vegetation cover. The second paper by Fernandez-Oto et al. [226] demonstrates that the enigmatic fairy circles of the Namibia desert can be stabilized by applying a strong nonlocal coupling which models the competition process between the plants. Numerical and analytical results obtained by the authors are in close agreement with the field observations. The last paper by Martinez-Garcia et al. [227] discusses the formation of vegetation patterns in arid and semi arid landscapes. They attribute this formation to the competition between plants and not to the facilitation phenomenon.

Acknowledgment

M.G.C. acknowledge the financial support of FONDECYT project 1120320. C.F.O. the financial support of Becas Chile. M.T received support from the Fonds National de la Recherche Scientifique (Belgium). This research was supported by the Interuniversity Attraction Poles program of the Belgian Science Policy Office, under grant IAP P7-35.

-
- [1] Glansdorff, P. & Prigogine, I., 1971 *Thermodynamic Theory of Structure, Stability and Fluctuations*. Wiley. ISBN 9780471302803.
 - [2] Prigogine, I. & Lefever, R., 1968 Symmetry breaking instabilities in dissipative systems. ii. *The Journal of Chemical Physics* **48**, 1695–1700. (doi:http://dx.doi.org/10.1063/1.1668896).
 - [3] Turing, A. M., 1952 The chemical basis of morphogenesis. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **237**,

- 37–72. (doi:10.1098/rstb.1952.0012).
- [4] BÃ©nard, H., 1901 *Les tourbillons cellulaires dans une nappe liquide propageant de la chaleur par convection, en rÃ©gime permanent*. Ph.D. thesis, UniversitÃ© de Paris. ThÃ©se de doctorat Sciences physiques FacultÃ© des sciences de l'UniversitÃ© de Paris 1901.
 - [5] Castets, V., Dulos, E., Boissonade, J. & De Kepper, P., 1990 Experimental evidence of a sustained standing turing-type nonequilibrium chemical pattern. *Phys. Rev. Lett.* **64**, 2953–2956. (doi:10.1103/PhysRevLett.64.2953).
 - [6] Ouyang, Q. & Swinney, H. L., 1991 Transition from a uniform state to hexagonal and striped turing patterns. *Nature* **352**, 610–612. (doi:10.1038/352610a0).
 - [7] Belousov, B. P., 1959 A periodic reaction and its mechanism. *Compilation of Abstracts on Radiation Medicine* **147**, 1.
 - [8] N., Z. A. & M., Z. A., 1970 Concentration wave propagation in two-dimensional liquid-phase self-oscillating system. *Nature* **225**, 535–537. (doi:10.1038/225535b0).
 - [9] Lefever, R. & Lejeune, O., 1997 On the origin of tiger bush. *Bulletin of Mathematical Biology* **59**, 263–294. ISSN 0092-8240. (doi:10.1007/BF02462004).
 - [10] Lugiato, L. A., 1994 Transverse nonlinear optics: Introduction and review. *Chaos, Solitons & Fractals* **4**, 1251. ISSN 0960-0779. (doi:http://dx.doi.org/10.1016/0960-0779(94)90080-9). Special Issue: Nonlinear Optical Structures, Patterns, Chaos/ce:title.
 - [11] Rosanov, N. N., 1996 I transverse patterns in wide-aperture nonlinear optical systems. In *Progress in Optics* (ed. E. Wolf), volume 35 of *Progress in Optics*, pp. 1 – 60. Elsevier. (doi:http://dx.doi.org/10.1016/S0079-6638(08)70527-4).
 - [12] Mandel, P., 1997 *Theoretical Problems in Cavity Nonlinear Optics*. Cambridge University Press. ISBN 9780511529337. Cambridge Books Online.
 - [13] Arecchi, F., Boccaletti, S. & Ramazza, P., 1999 Pattern formation and competition in nonlinear optics. *Physics Reports* **318**, 1 – 83. ISSN 0370-1573. (doi:http://dx.doi.org/10.1016/S0370-1573(99)00007-1).
 - [14] Pismen, L. M., 1999 *Vortices in Nonlinear Fields: From Liquid Crystals to Superfluids, from Non-equilibrium Patterns to Cosmic Strings*, volume 100 of *International series of monographs on physics*. Oxford University Press, clarendon press edition.
 - [15] Rosanov, N. N., 2002 *Spatial Hysteresis and Optical Patterns*. Springer Series in Synergetics. Springer Berlin Heidelberg. (doi:10.1007/978-3-662-04792-7).
 - [16] Staliunas, K. & SÃ¡nchez-Morcillo, V., 2003 *Transverse Patterns in Nonlinear Optical Resonators*. Springer tracts in modern physics. Springer Berlin Heidelberg. (doi:10.1007/3-540-36416-1).
 - [17] Kivshar, Y. S. & Agrawal, G. P., 2003 *Optical Solitons: From Fibers to Photonic Crystals*. Elsevier.
 - [18] Murray, J. D., 1989 *Mathematical Biology*, volume 19 of *Biomathematics*. Springer Berlin Heidelberg. (doi:10.1007/978-3-662-08539-4).
 - [19] Denz, C., Schwab, M. & Weilnau, C., 2003 *Transverse-Pattern Formation in Photorefractive Optics*, volume 188 of *Springer Tracts in Modern Physics*. Springer Berlin Heidelberg. (doi:10.1007/b13583).
 - [20] Mandel, P. & Tlidi, M., 2004 Transverse dynamics in cavity nonlinear optics (2000-2003). *Journal of Optics B: Quantum and Semiclassical Optics* **6**, R60.
 - [21] Malomed, B. A., Mihalache, D., Wise, F. & Torner, L., 2005 Spatiotemporal optical solitons. *Journal of Optics B: Quantum and Semiclassical Optics* **7**, R53.
 - [22] Mikhailov, A. S. & Showalter, K., 2006 Control of waves, patterns and turbulence in chemical systems. *Physics Reports* **425**, 79 – 194. ISSN 0370-1573. (doi:http://dx.doi.org/10.1016/j.physrep.2005.11.003).
 - [23] Tlidi, M., Taki, M. & Kolokolnikov, T., 2007 Introduction: dissipative localized structures in extended systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science* **17**, 037101.
 - [24] Akhmediev, N. & Ankiewicz, A. (eds.), 2008 *Dissipative Solitons: from Optics to Biology and Medicine*, volume 751 of *Lecture Notes in Physics*. Springer.
 - [25] Ackemann, T., Firth, W. J. & Oppo, G.-L., 2009 Fundamentals and applications of spatial dissipative solitons in photonic devices. volume 57 of *Advances In Atomic, Molecular, and Optical Physics*, chapter 6, p. 323. Academic Press. (doi:http://dx.doi.org/10.1016/S1049-250X(09)57006-1).
 - [26] Kuszelewicz, R., Barbay, S., Tissoni, G. & Almuneau, G., 2010 Editorial on dissipative optical solitons. *Eur. Phys. J. D* **59**, 1. (doi:10.1140/epjd/e2010-00167-7).
 - [27] Purwins, H. G., BÃ¼dcker, H. U. & Amiranashvili, S., 2010 Dissipative solitons. *Advances in Physics* **59**, 485. (doi:10.1080/00018732.2010.498228).
 - [28] Ridolfi, L., D'Odorico, P. & Laio, F., 2011 *Noise-Induced Phenomena in the Environmental Sciences*. cambridge University Press.
 - [29] Descalzi, O., Clerc, M. & Residori, S., 2011 *Localized States in Physics: Solitons and Patterns*. Springer. ISBN 9783642165498.
 - [30] Leblond, H. & Mihalache, D., 2013 Models of few optical cycle solitons beyond the slowly varying envelope approximation. *Physics Reports* **523**, 61. ISSN 0370-1573. (doi:http://dx.doi.org/10.1016/j.physrep.2012.10.006).
 - [31] Graham, R. & Haken, H., 1970 Laserlight - first example of a second-order phase transition far away from thermal equilibrium. *Zeitschrift fÃ¼r Physik* **237**, 31–46. ISSN 0044-3328. (doi:10.1007/BF01400474).
 - [32] Akhmanov, S., Khoklov, R. & Sukhorukov, A., 1972 Self-focusing, self-defocusing and self-modulation of laser beams. Number 2 in *Laser Handbook*, chapter E3, p. 1151. North Holland.
 - [33] McLaughlin, D. W., Moloney, J. V. & Newell, A. C., 1985 New class of instabilities in passive optical cavities. *Phys. Rev. Lett.* **54**, 681–684. (doi:10.1103/PhysRevLett.54.681).
 - [34] Lugiato, L. A., Oldano, C. & Narducci, L. M., 1988 Cooperative frequency locking and stationary spatial structures in lasers. *J. Opt. Soc. Am. B* **5**, 879–888. (doi:10.1364/JOSAB.5.000879).
 - [35] Couillet, P., Gil, L. & Rocca, F., 1989 Optical vortices. *Optics Communications* **73**, 403 – 408. ISSN 0030-4018. (doi:http://dx.doi.org/10.1016/0030-4018(89)90180-6).
 - [36] Brambilla, M., Battipede, F., Lugiato, L. A., Penna, V., Prati, F., Tamm, C. & Weiss, C. O., 1991 Transverse laser patterns. i. phase singularity crystals. *Phys. Rev. A* **43**, 5090–5113. (doi:10.1103/PhysRevA.43.5090).
 - [37] Oppo, G.-L., D'Alessandro, G. & Firth, W. J., 1991 Spatiotemporal instabilities of lasers in models reduced via center manifold techniques. *Phys. Rev. A* **44**, 4712–

4720. (doi:10.1103/PhysRevA.44.4712).
- [38] Staliunas, K., 1993 Laser ginzburg-landau equation and laser hydrodynamics. *Phys. Rev. A* **48**, 1573–1581. (doi:10.1103/PhysRevA.48.1573).
 - [39] Jakobsen, P. K., Moloney, J. V., Newell, A. C. & Indik, R., 1992 Space-time dynamics of wide-gain-section lasers. *Phys. Rev. A* **45**, 8129–8137. (doi:10.1103/PhysRevA.45.8129).
 - [40] Residori, S., Petrossian, A., Nagaya, T. & Clerc, M., 2004 Localized structures and their dynamics in a liquid crystal light valve with optical feedback. *Journal of Optics B: Quantum and Semiclassical Optics* **6**, S169.
 - [41] Clerc, M. G., Petrossian, A. & Residori, S., 2005 Bouncing localized structures in a liquid-crystal light-valve experiment. *Phys. Rev. E* **71**, 015205. (doi:10.1103/PhysRevE.71.015205).
 - [42] Kozyreff, G. & Tlidi, M., 2007 Nonvariational real swift-hohenberg equation for biological, chemical, and optical systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science* **17**, 037103. (doi:http://dx.doi.org/10.1063/1.2759436).
 - [43] Staliunas, K., Tarroja, M. F. H., Sleky, G., Weiss, C. O. & Dambly, L., 1995 Analogy between photorefractive oscillators and class-a lasers. *Phys. Rev. A* **51**, 4140–4151. (doi:10.1103/PhysRevA.51.4140).
 - [44] Longhi, S., 1996 Traveling-wave states and secondary instabilities in optical parametric oscillators. *Phys. Rev. A* **53**, 4488–4499. (doi:10.1103/PhysRevA.53.4488).
 - [45] de Valcárcel, G. J., Staliunas, K., Roldán, E. & Sánchez-Morcillo, V. J., 1996 Transverse patterns in degenerate optical parametric oscillation and degenerate four-wave mixing. *Phys. Rev. A* **54**, 1609–1624. (doi:10.1103/PhysRevA.54.1609).
 - [46] Tlidi, M., Georgiou, M. & Mandel, P., 1993 Transverse patterns in nascent optical bistability. *Phys. Rev. A* **48**, 4605–4609. (doi:10.1103/PhysRevA.48.4605).
 - [47] Staliunas, K., 1998 Three-dimensional turing structures and spatial solitons in optical parametric oscillators. *Phys. Rev. Lett.* **81**, 81–84. (doi:10.1103/PhysRevLett.81.81).
 - [48] Tlidi, M., Haelterman, M. & Mandel, P., 1998 Three-dimensional structures in diffractive and dispersive nonlinear ring cavities. *Quantum and Semiclassical Optics: Journal of the European Optical Society Part B* **10**, 869.
 - [49] Tlidi, M., 2000 Three-dimensional crystals and localized structures in diffractive and dispersive nonlinear ring cavities. *Journal of Optics B: Quantum and Semiclassical Optics* **2**, 438.
 - [50] Tassin, P., der Sande, G. V., Veretenov, N., Kockaert, P., Veretennicoff, I. & Tlidi, M., 2006 Three-dimensional structures in nonlinear cavities containing left-handed materials. *Opt. Express* **14**, 9338–9343. (doi:10.1364/OE.14.009338).
 - [51] Veretenov, N. & Tlidi, M., 2009 Dissipative light bullets in an optical parametric oscillator. *Phys. Rev. A* **80**, 023822. (doi:10.1103/PhysRevA.80.023822).
 - [52] Dai, C.-Q., Wang, X.-G. & Zhou, G.-Q., 2014 Stable light-bullet solutions in the harmonic and parity-time-symmetric potentials. *Phys. Rev. A* **89**, 013834. (doi:10.1103/PhysRevA.89.013834).
 - [53] Brambilla, M., Maggipinto, T., Patera, G. & Columbo, L., 2004 Cavity light bullets: Three-dimensional localized structures in a nonlinear optical resonator. *Phys. Rev. Lett.* **93**, 203901. (doi:10.1103/PhysRevLett.93.203901).
 - [54] Blaauwboer, M., Kurizki, G. & Malomed, B. A., 2000 Spatiotemporally localized solitons in resonantly absorbing bragg reflectors. *Phys. Rev. E* **62**, R57–R59. (doi:10.1103/PhysRevE.62.R57).
 - [55] de Valcárcel, G. J. & Staliunas, K., 2003 Excitation of phase patterns and spatial solitons via two-frequency forcing of a 1:1 resonance. *Phys. Rev. E* **67**, 026604. (doi:10.1103/PhysRevE.67.026604).
 - [56] de Valcárcel, G. J. & Staliunas, K., 2010 Pattern formation through phase bistability in oscillatory systems with space-modulated forcing. *Phys. Rev. Lett.* **105**, 054101. (doi:10.1103/PhysRevLett.105.054101).
 - [57] Fernandez-Oto, C., de Valcárcel, G. J., Tlidi, M., Panajotov, K. & Staliunas, K., 2014 Phase-bistable patterns and cavity solitons induced by spatially periodic injection into vertical-cavity surface-emitting lasers. *Phys. Rev. A* **89**, 055802. (doi:10.1103/PhysRevA.89.055802).
 - [58] Staliunas, K. & Tlidi, M., 2005 Hyperbolic transverse patterns in nonlinear optical resonators. *Phys. Rev. Lett.* **94**, 133902. (doi:10.1103/PhysRevLett.94.133902).
 - [59] Kolpakov, S., Esteban-Martín, A., Silva, F., García, J., Staliunas, K. & de Valcárcel, G. n. J., 2008 Experimental demonstration of hyperbolic patterns. *Phys. Rev. Lett.* **101**, 254101. (doi:10.1103/PhysRevLett.101.254101).
 - [60] Staliunas, K., 1992 Dynamics of optical vortices in a laser beam. *Optics Communications* **90**, 123 – 127. ISSN 0030-4018. (doi:http://dx.doi.org/10.1016/0030-4018(92)90342-O).
 - [61] Rozas, D., Law, C. T. & G. A. Swartzlander, J., 1997 Propagation dynamics of optical vortices. *J. Opt. Soc. Am. B* **14**, 3054–3065. (doi:10.1364/JOSAB.14.003054).
 - [62] Kivshar, Y. S. & Ostrovskaya, E. A., 2001 Optical vortices folding and twisting waves of light. *Opt. Photon. News* **12**, 24–28. (doi:10.1364/OPN.12.4.000024).
 - [63] Staliunas, K., 1994 Vortices and dark solitons in the two-dimensional nonlinear schrödinger equation. *Chaos, Solitons & Fractals* **4**, 1783 – 1796. ISSN 0960-0779. (doi:http://dx.doi.org/10.1016/0960-0779(94)90111-2). Special Issue: Nonlinear Optical Structures, Patterns, Chaos.
 - [64] Arecchi, F. T., Giacomelli, G., Ramazza, P. L. & Residori, S., 1991 Vortices and defect statistics in two-dimensional optical chaos. *Phys. Rev. Lett.* **67**, 3749–3752. (doi:10.1103/PhysRevLett.67.3749).
 - [65] Gil, L., Emilsson, K. & Oppo, G.-L., 1992 Dynamics of spiral waves in a spatially inhomogeneous hopf bifurcation. *Phys. Rev. A* **45**, R567–R570. (doi:10.1103/PhysRevA.45.R567).
 - [66] Staliunas, K. & Weiss, C., 1995 Tilted and standing waves and vortex lattices in class-a lasers. *Physica D: Nonlinear Phenomena* **81**, 79 – 93. ISSN 0167-2789. (doi:http://dx.doi.org/10.1016/0167-2789(94)00193-T).
 - [67] Hennequin, D., Dambly, L., Dangoisse, D. & Glorieux, P., 1994 Basic transverse dynamics of a photorefractive oscillator. *J. Opt. Soc. Am. B* **11**, 676–684. (doi:10.1364/JOSAB.11.000676).
 - [68] Staliunas, K. & Weiss, C. O., 1995 Nonstationary vortex lattices in large-aperture class b lasers. *J. Opt. Soc. Am.*

- B* **12**, 1142–1149. (doi:10.1364/JOSAB.12.001142).
- [69] Barboza, R., Bortolozzo, U., Assanto, G., Vidal-Henriquez, E., Clerc, M. G. & Residori, S., 2012 Vortex induction via anisotropy stabilized light-matter interaction. *Phys. Rev. Lett.* **109**, 143901. (doi:10.1103/PhysRevLett.109.143901).
- [70] Barboza, R., Bortolozzo, U., Assanto, G., Vidal-Henriquez, E., Clerc, M. G. & Residori, S., 2013 Harnessing optical vortex lattices in nematic liquid crystals. *Phys. Rev. Lett.* **111**, 093902. (doi:10.1103/PhysRevLett.111.093902).
- [71] Haelterman, M. & Sheppard, A., 1994 Polarization instability, multistability and transverse localized structures in kerr media. *Chaos, Solitons & Fractals* **4**, 1731 – 1743. ISSN 0960-0779. (doi:http://dx.doi.org/10.1016/0960-0779(94)90107-4). Special Issue: Nonlinear Optical Structures, Patterns, Chaos.
- [72] Kivshar, Y. S. & Yang, X., 1994 Dynamics of dark solitons. *Chaos, Solitons & Fractals* **4**, 1745 – 1758. ISSN 0960-0779. (doi:http://dx.doi.org/10.1016/0960-0779(94)90108-2). Special Issue: Nonlinear Optical Structures, Patterns, Chaos.
- [73] Swartzlander, G. A. & Law, C. T., 1992 Optical vortex solitons observed in kerr nonlinear media. *Phys. Rev. Lett.* **69**, 2503–2506. (doi:10.1103/PhysRevLett.69.2503).
- [74] Firth, W. J. & Paré, C., 1988 Transverse modulational instabilities for counterpropagating beams in kerr media. *Opt. Lett.* **13**, 1096–1098. (doi:10.1364/OL.13.001096).
- [75] Akhmanov, S. A., Vorontsov, M. A., Ivanov, V. Y., Larichev, A. V. & Zheleznykh, N. I., 1992 Controlling transverse-wave interactions in nonlinear optics: generation and interaction of spatiotemporal structures. *J. Opt. Soc. Am. B* **9**, 78–90. (doi:10.1364/JOSAB.9.000078).
- [76] Zheleznykh, N., Berre, M. L., Ressayre, E. & Tallet, A., 1994 Rotating spiral waves in a nonlinear optical system with spatial interactions. *Chaos, Solitons & Fractals* **4**, 1717 – 1728. ISSN 0960-0779. (doi:http://dx.doi.org/10.1016/0960-0779(94)90106-6). Special Issue: Nonlinear Optical Structures, Patterns, Chaos.
- [77] Fauve, S. & Thual, O., 1990 Solitary waves generated by subcritical instabilities in dissipative systems. *Phys. Rev. Lett.* **64**, 282–284. (doi:10.1103/PhysRevLett.64.282).
- [78] Sleky, G., Staliunas, K. & Weiss, C., 1998 Spatial localized structures in resonators with saturable absorber. *Optics Communications* **149**, 113 – 116. ISSN 0030-4018. (doi:http://dx.doi.org/10.1016/S0030-4018(97)00667-6).
- [79] Tlidi, M., Mandel, P. & Lefever, R., 1994 Localized structures and localized patterns in optical bistability. *Phys. Rev. Lett.* **73**, 640. (doi:10.1103/PhysRevLett.73.640).
- [80] Scroggie, A. J., Firth, W. J., McDonald, G. S., Tlidi, M., Lefever, R. & Lugiato, L. A., 1994 Pattern formation in a passive kerr cavity. *Chaos, Solitons and Fractals* **4**, 1323. ISSN 0960-0779. (doi:http://dx.doi.org/10.1016/0960-0779(94)90084-1).
- [81] Staliunas, K. & Sánchez-Morcillo, V. J., 1997 Localized structures in degenerate optical parametric oscillators. *Optics Communications* **139**, 306 – 312. ISSN 0030-4018. (doi:http://dx.doi.org/10.1016/S0030-4018(97)00109-0).
- [82] Longhi, S., 1998 Spatial solitary waves and patterns in type ii second-harmonic generation. *Opt. Lett.* **23**, 346–348. (doi:10.1364/OL.23.000346).
- [83] Staliunas, K. & Sánchez-Morcillo, V. J., 1998 Dynamics of phase domains in the swift-hohenberg equation. *Physics Letters A* **241**, 28 – 34. ISSN 0375-9601. (doi: http://dx.doi.org/10.1016/S0375-9601(98)00084-X).
- [84] Tlidi, M., Mandel, P. & Lefever, R., 1998 Kinetics of localized pattern formation in optical systems. *Phys. Rev. Lett.* **81**, 979–982. (doi:10.1103/PhysRevLett.81.979).
- [85] Tlidi, M., Mandel, P., Berre, M. L., Ressayre, E., Tallet, A. & Menza, L. D., 2000 Phase-separation dynamics of circular domain walls in the degenerate optical parametric oscillator. *Opt. Lett.* **25**, 487–489. (doi: 10.1364/OL.25.000487).
- [86] Izús, G., San Miguel, M. & Santagiustina, M., 2001 Phase-locked spatial domains and bloch domain walls in type-ii optical parametric oscillators. *Phys. Rev. E* **64**, 056231. (doi:10.1103/PhysRevE.64.056231).
- [87] Barbay, S., Kuszelewicz, R. & Tredicce, J. R., 2011 Cavity solitons in vcsel devices. *Advances in Optical Technologies* **2011**, 628761.
- [88] Soda, H., Iga, K., Kitahara, C. & Suematsu, Y., 1979 Gainasp/inp surface emitting injection lasers. *Japanese Journal of Applied Physics* **18**, 2329.
- [89] Li, H. E. & Iga, K. (eds.), 2003 *Vertical-Cavity Surface-Emitting Laser Devices*, volume 6 of *Springer Series in Photonics*. Springer Berlin Heidelberg.
- [90] Michalzick, R. (ed.), 2013 *VCELS: Fundamentals, Technology and Applications of Vertical-Cavity Surface-Emitting Lasers*, volume 166 of *Springer Series in Optical Sciences*. Springer Berlin Heidelberg.
- [91] Panajotov, K., Sciamanna, M., Arteaga, M. & Thienpont, H., 2013 Optical feedback in vertical-cavity surface-emitting lasers. *Selected Topics in Quantum Electronics, IEEE Journal of* **19**, 1700312–1700312. ISSN 1077-260X. (doi:10.1109/JSTQE.2012.2235060).
- [92] Panajotov, K., Gatare, I., Valle, A., Thienpont, H. & Sciamanna, M., 2009 Polarization- and transverse-mode dynamics in optically injected and gain-switched vertical-cavity surface-emitting lasers. *Quantum Electronics, IEEE Journal of* **45**, 1473–1481. ISSN 0018-9197. (doi:10.1109/JQE.2009.2024958).
- [93] Taranenko, V. B., Ganne, I., Kuszelewicz, R. J. & Weiss, C. O., 2000 Patterns and localized structures in bistable semiconductor resonators. *Phys. Rev. A* **61**, 063818. (doi:10.1103/PhysRevA.61.063818).
- [94] Averlant, E., Tlidi, M., Thienpont, H., Ackemann, T. & Panajotov, K., 2014 Experimental observation of localized structures in medium size vcsels. *Opt. Express* **22**, 762–772. (doi:10.1364/OE.22.000762).
- [95] Barland, S., Tredicce, J. R., Brambilla, M., Lugiato, L. A., Balle, S., Giudici, M., Maggipinto, T., Spinelli, L., Tissoni, G., Knödl, T. *et al.*, 2002 Cavity solitons as pixels in semiconductor microcavities. *Nature* **419**, 699.
- [96] Hachair, X., Pedaci, F., Caboche, E., Barland, S., Giudici, M., Tredicce, J. R., Prati, F., Tissoni, G., Kheradmand, R., Lugiato, L. A. *et al.*, 2006 Cavity solitons in a driven vcsel above threshold. *Selected Topics in Quantum Electronics, IEEE Journal of* **12**, 339–351. ISSN

- 1077-260X. (doi:10.1109/JSTQE.2006.872711).
- [97] Hachair, X., Tissoni, G., Thienpont, H. & Panajotov, K., 2009 Linearly polarized bistable localized structure in medium-size vertical-cavity surface-emitting lasers. *Phys. Rev. A* **79**, 011801. (doi:10.1103/PhysRevA.79.011801).
- [98] Tanguy, Y., Ackemann, T., Firth, W. J. & Jäger, R., 2008 Realization of a semiconductor-based cavity soliton laser. *Phys. Rev. Lett.* **100**, 013907. (doi:10.1103/PhysRevLett.100.013907).
- [99] Paulau, P. V., Gomila, D., Ackemann, T., Loiko, N. A. & Firth, W. J., 2008 Self-localized structures in vertical-cavity surface-emitting lasers with external feedback. *Phys. Rev. E* **78**, 016212. (doi:10.1103/PhysRevE.78.016212).
- [100] Genevet, P., Barland, S., Giudici, M. & Tredicce, J. R., 2008 Cavity soliton laser based on mutually coupled semiconductor microresonators. *Phys. Rev. Lett.* **101**, 123905. (doi:10.1103/PhysRevLett.101.123905).
- [101] Columbo, L., Prati, F., Brambilla, M. & Maggipinto, T., 2010 Self-pulsing localized structures in a laser with saturable absorber. *The European Physical Journal D* **59**, 115–120. ISSN 1434-6060. (doi:10.1140/epjd/e2010-00110-0).
- [102] Rosanov, N. N. & Khodova, G. V., 1988 Autosolitons in bistable interferometers. *Optics and Spectroscopy* **65**, 449–450.
- [103] Vladimirov, A. G., Fedorov, S. V., Kaliteevskii, N. A., Khodova, G. V. & Rosanov, N. N., 1999 Numerical investigation of laser localized structures. *Journal of Optics B: Quantum and Semiclassical Optics* **1**, 101.
- [104] Fedorov, S. V., Vladimirov, A. G., Khodova, G. V. & Rosanov, N. N., 2000 Effect of frequency detunings and finite relaxation rates on laser localized structures. *Phys. Rev. E* **61**, 5814–5824. (doi:10.1103/PhysRevE.61.5814).
- [105] Elsass, T., Gauthron, K., Beaudoin, G., Sagnes, I., Kuszelewicz, R. & Barbay, S., 2010 Control of cavity solitons and dynamical states in a monolithic vertical cavity laser with saturable absorber. *The European Physical Journal D* **59**, 91–96. ISSN 1434-6060. (doi:10.1140/epjd/e2010-00079-6).
- [106] Elsass, T., Gauthron, K., Beaudoin, G., Sagnes, I., Kuszelewicz, R. & Barbay, S., 2010 Fast manipulation of laser localized structures in a monolithic vertical cavity with saturable absorber. *Applied Physics B* **98**, 327–331. ISSN 0946-2171. (doi:10.1007/s00340-009-3748-9).
- [107] Pedaci, F., Genevet, P., Barland, S., Giudici, M. & Tredicce, J. R., 2006 Positioning cavity solitons with a phase mask. *Applied Physics Letters* **89**, 221111. (doi: http://dx.doi.org/10.1063/1.2388867).
- [108] Pedaci, F., Barland, S., Caboche, E., Genevet, P., Giudici, M., Tredicce, J. R., Ackemann, T., Scroggie, A. J., Firth, W. J., Oppo, G. L. *et al.*, 2008 All-optical delay line using semiconductor cavity solitons. *Appl. Phys. Lett.* **92**, 011101. (doi:10.1063/1.2828458).
- [109] Pedaci, F., Tissoni, G., Barland, S., Giudici, M. & Tredicce, J., 2008 Mapping local defects of extended media using localized structures. *Appl. Phys. Lett.* **93**, 111104. (doi:10.1063/1.2977603).
- [110] Staliunas, K., 2003 Midband dissipative spatial solitons. *Phys. Rev. Lett.* **91**, 053901. (doi:10.1103/PhysRevLett.91.053901).
- [111] Staliunas, K., Egorov, O., Kivshar, Y. S. & Lederer, F., 2008 Bloch cavity solitons in nonlinear resonators with intracavity photonic crystals. *Phys. Rev. Lett.* **101**, 153903. (doi:10.1103/PhysRevLett.101.153903).
- [112] Vladimirov, A. G., Skryabin, D. V., Kozyreff, G., Mandel, P. & Tlidi, M., 2006 Bragg localized structures in a passive cavity with transverse modulation of the refractive index and the pump. *Opt. Express* **14**, 1–6. (doi:10.1364/OPEX.14.000001).
- [113] Kasprzak, J., Richard, M., Kundermann, S., Baas, A., Jeambrun, P., Keeling, J. M. J., Marchetti, F. M., Szymanska, M. H., Andre, R., Staehli, J. L. *et al.*, 2006 Bose-einstein condensation of exciton polaritons. *Nature* **443**, 409–414. (doi:10.1038/nature05131).
- [114] Egorov, O. A., Skryabin, D. V., Yulin, A. V. & Lederer, F., 2009 Bright cavity polariton solitons. *Phys. Rev. Lett.* **102**, 153904. (doi:10.1103/PhysRevLett.102.153904).
- [115] Gibbs, H. M., Khitrova, G. & Koch, S. W., 2011 Exciton-polariton light-semiconductor coupling effects. *Nature Photonics* **5**, 273. (doi:10.1038/nphoton.2011.15).
- [116] Veselago, V. G., 1968 The electrodynamics of substances with simultaneously negative values of ϵ and μ . *Soviet Physics Uspekhi* **10**, 509.
- [117] Shelby, R. A., Smith, D. R. & Schultz, S., 2001 Experimental verification of a negative index of refraction. *Science* **292**, 77–79. (doi:10.1126/science.1058847).
- [118] Wiltshire, M. C. K., 2001 Bending light the wrong way. *Science* **292**, 60–61. (doi:10.1126/science.1060221).
- [119] Shalaev, V. M., 2007 Optical negative-index metamaterials. *Nat Photon* **1**, 41. (doi:10.1038/nphoton.2006.49).
- [120] Boardman, A., Brongersma, M., Stockman, M. & Wegener, M., 2009 Plasmonics and metamaterials: Introduction. *J. Opt. Soc. Am. B* **26**, PM1–PM1. (doi:10.1364/JOSAB.26.000PM1).
- [121] Kockaert, P., Tassin, P., Van der Sande, G., Veretennicoff, I. & Tlidi, M., 2006 Negative diffraction pattern dynamics in nonlinear cavities with left-handed materials. *Phys. Rev. A* **74**, 033822. (doi:10.1103/PhysRevA.74.033822).
- [122] Kockaert, P., Tassin, P., Veretennicoff, I., der Sande, G. V. & Tlidi, M., 2009 Beyond the zero-diffraction regime in optical cavities with a left-handed material. *J. Opt. Soc. Am. B* **26**, B148–B155. (doi:10.1364/JOSAB.26.00B148).
- [123] Tassin, P., Gelens, L., Danckaert, J., Veretennicoff, I., Van der Sande, G., Kockaert, P. & Tlidi, M., 2007 Dissipative structures in left-handed material cavity optics. *Chaos: An Interdisciplinary Journal of Nonlinear Science* **17**, 037116. (doi:http://dx.doi.org/10.1063/1.2768158).
- [124] Gelens, L., Van der Sande, G., Tassin, P., Tlidi, M., Kockaert, P., Gomila, D., Veretennicoff, I. & Danckaert, J., 2007 Impact of nonlocal interactions in dissipative systems: Towards minimal-sized localized structures. *Phys. Rev. A* **75**, 063812. (doi:10.1103/PhysRevA.75.063812).
- [125] Kozyrev, A. B., Shadrivov, I. V. & Kivshar, Y. S., 2014 Soliton generation in active nonlinear metamaterials. *Applied Physics Letters* **104**, 084105. (doi: http://dx.doi.org/10.1063/1.4866856).
- [126] Pomeau, Y., 1986 Front motion, metastability and subcritical bifurcations in hydrodynamics. *Physica D: Nonlinear Phenomena* **23**, 3 – 11. ISSN 0167-2789. (doi:

- [http://dx.doi.org/10.1016/0167-2789\(86\)90104-1](http://dx.doi.org/10.1016/0167-2789(86)90104-1).
- [127] Langer, J. S., 1980 Instabilities and pattern formation in crystal growth. *Rev. Mod. Phys.* **52**, 1–28. (doi:10.1103/RevModPhys.52.1).
 - [128] Collet, P. & Eckman, J.-P., 2014 *Instabilities and Fronts in Extended Systems*. Princeton Series in Physics. Princeton University Press.
 - [129] Eschenfelder, A. H., 1980 *Magnetic Bubble Technology*, volume 14 of *Springer Series in Solid-State Sciences*. Springer Berlin Heidelberg. (doi:10.1007/978-3-642-96549-4).
 - [130] Börzsönyi, T., Akamatsu, S. & Faivre, G., 2009 Weakly faceted cellular patterns versus growth-induced plastic deformation in thin-sample directional solidification of monoclinic biphenyl. *Phys. Rev. E* **80**, 051601. (doi:10.1103/PhysRevE.80.051601).
 - [131] Clerc, M. G., Residori, S. & Riera, C. S., 2001 First-order fr edericksz transition in the presence of light-driven feedback in nematic liquid crystals. *Phys. Rev. E* **63**, 060701. (doi:10.1103/PhysRevE.63.060701).
 - [132] Gomila, D., Colet, P., Oppo, G.-L. & San Miguel, M., 2001 Stable droplets and growth laws close to the modulational instability of a domain wall. *Phys. Rev. Lett.* **87**, 194101. (doi:10.1103/PhysRevLett.87.194101).
 - [133] Clerc, M., Nagaya, T., Petrossian, A., Residori, S. & Riera, C., 2004 First-order fr edericksz transition and front propagation in a liquid crystal light valve with feedback. *The European Physical Journal D - Atomic, Molecular, Optical and Plasma Physics* **28**, 435–445. ISSN 1434-6060. (doi:10.1140/epjd/e2003-00316-1).
 - [134] Residori, S., 2005 Patterns, fronts and structures in a liquid-crystal-light-valve with optical feedback. *Physics Reports* **416**, 201 – 272. ISSN 0370-1573. (doi:http://dx.doi.org/10.1016/j.physrep.2005.06.004).
 - [135] Petrov, V., Ouyang, Q. & Swinney, H. L., 1997 Resonant pattern formation in a chemical system. *Nature* **388**, 655.
 - [136] Aranson, I. & Tsimring, L., 2008 *Granular Patterns*. Oxford University Press. (doi:10.1093/acprof:oso/9780199534418.001.0001).
 - [137] Melo, F., Umbanhowar, P. B. & Swinney, H. L., 1995 Hexagons, kinks, and disorder in oscillated granular layers. *Phys. Rev. Lett.* **75**, 3838–3841. (doi:10.1103/PhysRevLett.75.3838).
 - [138] Douady, S., Fauve, S. & Laroche, C., 1989 Subharmonic instabilities and defects in a granular layer under vertical vibrations. *EPL (Europhysics Letters)* **8**, 621.
 - [139] Moon, S. J., Shattuck, M. D., Bizon, C., Goldman, D. I., Swift, J. B. & Swinney, H. L., 2001 Phase bubbles and spatiotemporal chaos in granular patterns. *Phys. Rev. E* **65**, 011301. (doi:10.1103/PhysRevE.65.011301).
 - [140] Moon, S. J., Goldman, D. I., Swift, J. B. & Swinney, H. L., 2003 Kink-induced transport and segregation in oscillated granular layers. *Phys. Rev. Lett.* **91**, 134301. (doi:10.1103/PhysRevLett.91.134301).
 - [141] Mac as, J. E., Clerc, M. G., Falcon, C. & Garcia-Nustes, M. A., 2013 Spatially modulated kinks in shallow granular layers. *Phys. Rev. E* **88**, 020201. (doi:10.1103/PhysRevE.88.020201).
 - [142] Fisher, R. A., 1937 The wave of advance of advantageous genes. *Annals of Eugenics* **7**, 355–369. ISSN 2050-1439. (doi:10.1111/j.1469-1809.1937.tb02153.x).
 - [143] Clerc, M. G., Escaff, D. & Kenkre, V. M., 2005 Patterns and localized structures in population dynamics. *Phys. Rev. E* **72**, 056217. (doi:10.1103/PhysRevE.72.056217).
 - [144] van Saarloos, W. & Hohenberg, P., 1992 Fronts, pulses, sources and sinks in generalized complex ginzburg-landau equations. *Physica D: Nonlinear Phenomena* **56**, 303 – 367. ISSN 0167-2789. (doi:http://dx.doi.org/10.1016/0167-2789(92)90175-M).
 - [145] Coulet, P., 2002 Localized patterns and fronts in nonequilibrium systems. *International Journal of Bifurcation and Chaos* **12**, 2445–2457. (doi:10.1142/S021812740200614X).
 - [146] Kolmogorov, A., Petrovsky, I. & Piskunov, N., 1937 Study of the diffusion equation coupled to an increase in mass and its application to a problem in biology. *Bull Uni Moskou Ser Int A1* **6**, 1–26.
 - [147] van Saarloos, W., 2003 Front propagation into unstable states. *Physics Reports* **386**, 29 – 222. ISSN 0370-1573. (doi:http://dx.doi.org/10.1016/j.physrep.2003.08.001).
 - [148] Benguria, R. D. & Depassier, M. C., 2002 Linear and nonlinear marginal stability for fronts of hyperbolic reaction diffusion equations. *Phys. Rev. E* **66**, 026607. (doi:10.1103/PhysRevE.66.026607).
 - [149] Ahlers, G. & Cannell, D. S., 1983 Vortex-front propagation in rotating couette-taylor flow. *Phys. Rev. Lett.* **50**, 1583–1586. (doi:10.1103/PhysRevLett.50.1583).
 - [150] Fineberg, J. & Steinberg, V., 1987 Vortex-front propagation in rayleigh-b nard convection. *Phys. Rev. Lett.* **58**, 1332–1335. (doi:10.1103/PhysRevLett.58.1332).
 - [151] Powers, T. R. & Goldstein, R. E., 1997 Pearling and pinching: Propagation of rayleigh instabilities. *Phys. Rev. Lett.* **78**, 2555–2558. (doi:10.1103/PhysRevLett.78.2555).
 - [152] Langer, J., 1992 An introduction to the kinetics of first-order phase transition. In *Solids Far from Equilibrium* (ed. C. Godr sche), Collection Alea-Saclay: Monographs and Texts in Statistical Physics, chapter 3, pp. 297–364. Cambridge University Press.
 - [153] Residori, S., Petrossian, A., Nagaya, T., Riera, C. & Clerc, M., 2004 Fronts and localized structures in a liquid-crystal-light-valve with optical feedback. *Physica D: Nonlinear Phenomena* **199**, 149 – 165. ISSN 0167-2789. (doi:http://dx.doi.org/10.1016/j.physd.2004.08.010). Trends in Pattern Formation: Stability , Control and Fluctuations.
 - [154] Goldstein, R. E., Gunaratne, G. H., Gil, L. & Coulet, P., 1991 Hydrodynamic and interfacial patterns with broken space-time symmetry. *Phys. Rev. A* **43**, 6700–6721. (doi:10.1103/PhysRevA.43.6700).
 - [155] Coulet, P., Lega, J., Houchmanzadeh, B. & Lajzerowicz, J., 1990 Breaking chirality in nonequilibrium systems. *Phys. Rev. Lett.* **65**, 1352–1355. (doi:10.1103/PhysRevLett.65.1352).
 - [156] Bulaevskii, L. N. & Ginzburg, V. L., 1964 Temperature dependence of the shape of the domain wall in ferromagnetics and ferroelectrics. *Soviet Physics JETP* **18**, 530.
 - [157] Clerc, M. G., Coulibaly, S. & Laroze, D., 2009 Nonvariational ising-bloch transition in parametrically driven systems. *International Journal of Bifurcation and Chaos* **19**, 2717–2726. (doi:10.1142/S0218127409024438).
 - [158] J.M. Gilli, M. Morabito & T. Frisch, 1994 Ising-bloch transition in a nematic liquid crystal. *J. Phys. II France* **4**, 319–331. (doi:10.1051/jp2:1994131).

- [159] Kawagishi, T., Mizuguchi, T. & Sano, M., 1995 Points, walls, and loops in resonant oscillatory media. *Phys. Rev. Lett.* **75**, 3768–3771. (doi:10.1103/PhysRevLett.75.3768).
- [160] Haim, D., Li, G., Ouyang, Q., McCormick, W. D., Swinney, H. L., Hagberg, A. & Meron, E., 1996 Breathing spots in a reaction-diffusion system. *Phys. Rev. Lett.* **77**, 190–193. (doi:10.1103/PhysRevLett.77.190).
- [161] Esteban-Martí n, A., Taranenko, V. B., García, J., de Valcárcel, G. n. J. & Roldán, E., 2005 Controlled observation of a nonequilibrium ising-bloch transition in a nonlinear optical cavity. *Phys. Rev. Lett.* **94**, 223903. (doi:10.1103/PhysRevLett.94.223903).
- [162] Bensimon, D., Shraiman, B. I. & Croquette, V., 1988 Nonadiabatic effects in convection. *Phys. Rev. A* **38**, 5461–5464. (doi:10.1103/PhysRevA.38.5461).
- [163] Clerc, M. G., Falcon, C. & Tirapegui, E., 2005 Additive noise induces front propagation. *Phys. Rev. Lett.* **94**, 148302. (doi:10.1103/PhysRevLett.94.148302).
- [164] Epstein, T. & Fineberg, J., 2004 Control of spatiotemporal disorder in parametrically excited surface waves. *Phys. Rev. Lett.* **92**, 244502. (doi:10.1103/PhysRevLett.92.244502).
- [165] Schwartz, M. E. & Solomon, T. H., 2008 Chemical reaction fronts in ordered and disordered cellular flows with opposing winds. *Phys. Rev. Lett.* **100**, 028302. (doi:10.1103/PhysRevLett.100.028302).
- [166] Thiam, A. R., Bremond, N. & Bibette, J., 2009 Breaking of an emulsion under an ac electric field. *Phys. Rev. Lett.* **102**, 188304. (doi:10.1103/PhysRevLett.102.188304).
- [167] Sbragaglia, M., Peters, A. M., Pirat, C., Borkent, B. M., Lammertink, R. G. H., Wessling, M. & Lohse, D., 2007 Spontaneous breakdown of superhydrophobicity. *Phys. Rev. Lett.* **99**, 156001. (doi:10.1103/PhysRevLett.99.156001).
- [168] Douarche, C., Buguin, A., Salman, H. & Libchaber, A., 2009 E. coli and oxygen: A motility transition. *Phys. Rev. Lett.* **102**, 198101. (doi:10.1103/PhysRevLett.102.198101).
- [169] Douglas, J. F., Efimenko, K., Fischer, D. A., Phelan, F. R. & Genzer, J., 2007 Propagating waves of self-assembly in organosilane monolayers. *Proceedings of the National Academy of Sciences* **104**, 10324–10329. (doi:10.1073/pnas.0703620104).
- [170] Shnerb, N. M., Louzoun, Y., Bettelheim, E. & Solomon, S., 2000 The importance of being discrete: Life always wins on the surface. *Proceedings of the National Academy of Sciences* **97**, 10322–10324. (doi:10.1073/pnas.180263697).
- [171] Armero, J., Lacasta, A. M., Ramáñez-Piscina, L., Casademunt, J., Sancho, J. M. & Saguá, F., 1996 Pattern and velocity selection of fronts propagating in modulated media. *EPL (Europhysics Letters)* **33**, 429.
- [172] Haudin, F., Elías, R. G., Rojas, R. G., Bortolozzo, U., Clerc, M. G. & Residori, S., 2009 Driven front propagation in 1d spatially periodic media. *Phys. Rev. Lett.* **103**, 128003. (doi:10.1103/PhysRevLett.103.128003).
- [173] Mitschke, F., Steinmeyer, G. & Schwache, A., 1996 Generation of one-dimensional optical turbulence. *Physica D: Nonlinear Phenomena* **96**, 251 – 258. ISSN 0167-2789. (doi:http://dx.doi.org/10.1016/0167-2789(96)00025-5). Measures of Spatio-Temporal Dynamics.
- [174] Coen, S. & Haelterman, M., 1997 Modulational instability induced by cavity boundary conditions in a normally dispersive optical fiber. *Phys. Rev. Lett.* **79**, 4139–4142. (doi:10.1103/PhysRevLett.79.4139).
- [175] Stratmann, M., Pagel, T. & Mitschke, F., 2005 Experimental observation of temporal soliton molecules. *Phys. Rev. Lett.* **95**, 143902. (doi:10.1103/PhysRevLett.95.143902).
- [176] Lugiato, L. A. & Lefever, R., 1987 Spatial dissipative structures in passive optical systems. *Phys. Rev. Lett.* **58**, 2209–2211. (doi:10.1103/PhysRevLett.58.2209).
- [177] Leo, F., Coen, S., Kockaert, P., Gorza, S.-P., Emplit, P. & Haelterman, M., 2010 Temporal cavity solitons in one-dimensional kerr media as bits in an all-optical buffer. *Nat Photon* **4**, 471. (doi:10.1038/nphoton.2010.120).
- [178] Coen, S., Tlidi, M., Emplit, P. & Haelterman, M., 1999 Convection versus dispersion in optical bistability. *Phys. Rev. Lett.* **83**, 2328–2331. (doi:10.1103/PhysRevLett.83.2328).
- [179] Coulibaly, S., Taki, M. & Tlidi, M., 2014 Universal power law for front propagation in all fiber resonators. *Opt. Express* **22**, 483–489. (doi:10.1364/OE.22.000483).
- [180] Cavalcanti, S. B., Cressoni, J. C., da Cruz, H. R. & Gouveia-Neto, A. S., 1991 Modulation instability in the region of minimum group-velocity dispersion of single-mode optical fibers via an extended nonlinear schrödinger equation. *Phys. Rev. A* **43**, 6162–6165. (doi:10.1103/PhysRevA.43.6162).
- [181] Pitois, S. & Millot, G., 2003 Experimental observation of a new modulational instability spectral window induced by fourth-order dispersion in a normally dispersive single-mode optical fiber. *Optics Communications* **226**, 415 – 422. ISSN 0030-4018. (doi: http://dx.doi.org/10.1016/j.optcom.2003.09.001).
- [182] Joly, N. Y., Omenetto, F. G., Efimov, A., Taylor, A. J., Knight, J. C. & Russell, P. S. J., 2005 Competition between spectral splitting and raman frequency shift in negative-dispersion slope photonic crystal fiber. *Optics Communications* **248**, 281 – 285. ISSN 0030-4018. (doi: http://dx.doi.org/10.1016/j.optcom.2004.11.091).
- [183] Russell, P. S., 2006 Photonic-crystal fibers. *J. Lightwave Technol.* **24**, 4729–4749.
- [184] Schmidberger, M., Chang, W., Russell, P. S. J. & Joly, N. Y., 2012 Influence of timing jitter on nonlinear dynamics of a photonic crystal fiber ring cavity. *Opt. Lett.* **37**, 3576–3578. (doi:10.1364/OL.37.003576).
- [185] Tlidi, M., Mussot, A., Louvergneaux, E., Kozyreff, G., Vladimirov, A. G. & Taki, M., 2007 Control and removal of modulational instabilities in low-dispersion photonic crystal fiber cavities. *Opt. Lett.* **32**, 662–664. (doi:10.1364/OL.32.000662).
- [186] Tlidi, M. & Gelens, L., 2010 High-order dispersion stabilizes dark dissipative solitons in all-fiber cavities. *Opt. Lett.* **35**, 306–308. (doi:10.1364/OL.35.000306).
- [187] Tlidi, M., Bahloul, L., Cherbi, L., Hariz, A. & Coulibaly, S., 2013 Drift of dark cavity solitons in a photonic-crystal fiber resonator. *Phys. Rev. A* **88**, 035802. (doi:10.1103/PhysRevA.88.035802).
- [188] Macfadyen, W. A., 1950 Vegetation patterns in the semi-desert plains of british somaliland. *The Geographical Journal* **116**, pp. 199–211. ISSN 00167398.
- [189] White, L. P., 1971 Vegetation stripes on sheet wash surfaces. *Journal of Ecology* **59**, pp. 615–622. ISSN

- 00220477.
- [190] Lefever, R. & Turner, J. W., 2012 A quantitative theory of vegetation patterns based on plant structure and the non-local f-kpp equation. *Comptes Rendus MATHématique* **340**, 818 – 828. ISSN 1631-0721. (doi: <http://dx.doi.org/10.1016/j.crme.2012.10.030>). Out of Equilibrium Dynamics.
- [191] Lejeune, O. & Tlidi, M., 1999 A model for the explanation of vegetation stripes (tiger bush). *Journal of Vegetation Science* **10**, 201–208. ISSN 1654-1103. (doi: [10.2307/3237141](https://doi.org/10.2307/3237141)).
- [192] Lefever, R., Barbier, N., Couteron, P. & Lejeune, O., 2009 Deeply gapped vegetation patterns: On crown/root allometry, criticality and desertification. *Journal of Theoretical Biology* **261**, 194 – 209. ISSN 0022-5193. (doi: <http://dx.doi.org/10.1016/j.jtbi.2009.07.030>).
- [193] Tlidi, M., Lefever, R. & Vladimirov, A., 2008 On vegetation clustering, localized bare soil spots and fairy circles. In *Dissipative Solitons: From Optics to Biology and Medicine*, volume 751 of *Lecture Notes in Physics*, pp. 1–22. Springer Berlin Heidelberg. ISBN 978-3-540-78216-2. (doi: [10.1007/978-3-540-78217-9_15](https://doi.org/10.1007/978-3-540-78217-9_15)).
- [194] Klausmeier, C. A., 1999 Regular and irregular patterns in semiarid vegetation. *Science* **284**, 1826–1828. (doi: [10.1126/science.284.5421.1826](https://doi.org/10.1126/science.284.5421.1826)).
- [195] Sherratt, J. A., 2005 An analysis of vegetation stripe formation in semi-arid landscapes. *Journal of Mathematical Biology* **51**, 183–197. ISSN 0303-6812. (doi: [10.1007/s00285-005-0319-5](https://doi.org/10.1007/s00285-005-0319-5)).
- [196] HilleRisLambers, R., Rietkerk, M., van den Bosch, F., Prins, H. H. T. & de Kroon, H., 2001 Vegetation pattern formation in semi-arid grazing systems. *Ecology* **82**, 50. (doi: [10.1890/0012-9658\(2001\)082\[0050:VPFISA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[0050:VPFISA]2.0.CO;2)).
- [197] Okayasu, T. & Aizawa, Y., 2001 Systematic analysis of periodic vegetation patterns. *Progress of Theoretical Physics* **106**, 705–720. (doi: [10.1143/PTP.106.705](https://doi.org/10.1143/PTP.106.705)).
- [198] Gilad, E., von Hardenberg, J., Provenzale, A., Shachak, M. & Meron, E., 2004 Ecosystem engineers: From pattern formation to habitat creation. *Phys. Rev. Lett.* **93**, 098105. (doi: [10.1103/PhysRevLett.93.098105](https://doi.org/10.1103/PhysRevLett.93.098105)).
- [199] Gowda, K., Riecke, H. & Silber, M., 2014 Transitions between patterned states in vegetation models for semi-arid ecosystems. *Phys. Rev. E* **89**, 022701. (doi: [10.1103/PhysRevE.89.022701](https://doi.org/10.1103/PhysRevE.89.022701)).
- [200] D’Odorico, P., Laio, F. & Ridolfi, L., 2006 Patterns as indicators of productivity enhancement by facilitation and competition in dryland vegetation. *Journal of Geophysical Research: Biogeosciences* **111**. ISSN 2156-2202. (doi: [10.1029/2006JG000176](https://doi.org/10.1029/2006JG000176)).
- [201] Lejeune, O., Tlidi, M. & Couteron, P., 2002 Localized vegetation patches: A self-organized response to resource scarcity. *Phys. Rev. E* **66**, 010901. (doi: [10.1103/PhysRevE.66.010901](https://doi.org/10.1103/PhysRevE.66.010901)).
- [202] Rietkerk, M., Dekker, S. C., de Ruiter, P. C. & van de Koppel, J., 2004 Self-organized patchiness and catastrophic shifts in ecosystems. *Science* **305**, 1926–1929. (doi: [10.1126/science.1101867](https://doi.org/10.1126/science.1101867)).
- [203] von Hardenberg, J., Meron, E., Shachak, M. & Zarmi, Y., 2001 Diversity of vegetation patterns and desertification. *Phys. Rev. Lett.* **87**, 198101. (doi: [10.1103/PhysRevLett.87.198101](https://doi.org/10.1103/PhysRevLett.87.198101)).
- [204] van Rooyen, M., Theron, G., van Rooyen, N., Jankowitz, W. & Matthews, W., 2004 Mysterious circles in the namib desert: review of hypotheses on their origin. *Journal of Arid Environments* **57**, 467 – 485. ISSN 0140-1963. (doi: [http://dx.doi.org/10.1016/S0140-1963\(03\)00111-3](http://dx.doi.org/10.1016/S0140-1963(03)00111-3)).
- [205] Jankowitz, W., Rooyen, M. V., Shaw, D., Kaumba, J. & Rooyen, N. V., 2008 Mysterious circles in the namib desert. *South African Journal of Botany* **74**, 332 – 334. ISSN 0254-6299. (doi: <http://dx.doi.org/10.1016/j.sajb.2007.10.010>).
- [206] Tschinkel, W. R., 2012 The life cycle and life span of namibian fairy circles. *PLoS ONE* **7**, e38056. (doi: [10.1371/journal.pone.0038056](https://doi.org/10.1371/journal.pone.0038056)).
- [207] Cramer, M. D. & Barger, N. N., 2013 Are namibian “fairy circles” the consequence of self-organizing spatial vegetation patterning? *PLoS ONE* **8**, e70876. (doi: [10.1371/journal.pone.0070876](https://doi.org/10.1371/journal.pone.0070876)).
- [208] Juergens, N., 2013 The biological underpinnings of namib desert fairy circles. *Science* **339**, 1618–1621. (doi: [10.1126/science.1222999](https://doi.org/10.1126/science.1222999)).
- [209] Getzin, S., Wiegand, K., Wiegand, T., Yizhaq, H., von Hardenberg, J. & Meron, E., 2014 Adopting a spatially explicit perspective to study the mysterious fairy circles of namibia. *Ecography* ISSN 1600-0587. (doi: [10.1111/ecog.00911](https://doi.org/10.1111/ecog.00911)).
- [210] Vladimirov, A., Pimenov, A., Gurevich, S., Panajotov, K., Averlant, E. & Tlidi, M., 2014 Cavity solitons in vertical-cavity surface-emitting lasers. *Philos. Trans. R. Soc. Lond. A* .
- [211] Turconi, M., Giudici, M. & Barland, S., 2014 Response of laser localized structures to external perturbations in coupled semiconductor microcavities. *Philos. Trans. R. Soc. Lond. A* .
- [212] Vahed, H., Prati, F., Turconi, M., Barland, S. & Tissoni, G., 2014 Periodic and chaotic localized structures in a semiconductor laser with saturable absorber. *Philos. Trans. R. Soc. Lond. A* .
- [213] Barboza, R., Bortolozzo, U., Clerc, M., Residori, S. & Vidal-Henriquez, E., 2014 Light-matter interaction induces a single positive vortex with swirling arms. *Philos. Trans. R. Soc. Lond. A* .
- [214] Noskov, R., Smirnova, D. & Kivshar, Y., 2014 Plasmonic kinks and walking solitons in nonlinear lattices of metal nanoparticles. *Philos. Trans. R. Soc. Lond. A* .
- [215] Ding, E., Tang, A. Y. S., Chow, K. W. & Malomed, B., 2014 Pinned modes in two-dimensional lossy lattices with local gain and nonlinearity. *Philos. Trans. R. Soc. Lond. A* .
- [216] Malomed, Y. H. B. A. & Mihalache, D., 2014 Localized modes in dissipative lattice media: An overview. *Philos. Trans. R. Soc. Lond. A* .
- [217] Rosanov, N., Sochilin, G., Vinokurova, V. & Vysotina, N., 2014 Spatial and temporal structures in cavities with oscillating boundaries. *Philos. Trans. R. Soc. Lond. A* .
- [218] de Valcárcel, G., Martínez-Quesada, M. & Staliunas, K., 2014 Phase-bistable pattern formation in oscillatory systems via rocking: Application to nonlinear optical systems. *Philos. Trans. R. Soc. Lond. A* .
- [219] Verschuere, N., Bortolozzo, U., Clerc, M. & Residori, S., 2014 Chaoticon: localised pattern with permanent dynamics. *Philos. Trans. R. Soc. Lond. A* .
- [220] Tesio, E., Robb, G. R. M., Oppo, G.-L., Gomes, P. M., Ackemann, T., Labeyrie, G., Kaiser, R. & Firth, W. J.,

- 2014 Self-organisation in cold atomic gases: a synchronisation perspective. *Philos. Trans. R. Soc. Lond. A* .
- [221] Sergeyev, S. V., 2014 Fast and slowly evolving vector solitons in mode locked fibre lasers. *Philos. Trans. R. Soc. Lond. A* .
- [222] Bahloul, L., Cherbi, L., Hariz, A. & Tlidi, M., 2014 Temporal localized structures in photonic crystal fiber resonators and their spontaneous symmetry breaking instability. *Philos. Trans. R. Soc. Lond. A* .
- [223] Chabchoub, A., Kibler, B., Dudley, J. M. & Akhmediev, N., 2014 Hydrodynamics of periodic breathers. *Philos. Trans. R. Soc. Lond. A* .
- [224] Gurevich, S., 2014 Time-delayed feedback control of breathing localized structures in a three-component reaction-diffusion system. *Philos. Trans. R. Soc. Lond. A* .
- [225] Couteron, P., Anthelme, F., Clerc, M., Escaff, D., Fernandez-Oto, C. & M.Tlidi, 2014 Clonal plant morphologies and spatial patterns as self-organized responses to resource-limited environments. *Philos. Trans. R. Soc. Lond. A* .
- [226] Fernandez-Oto, C., Tlidi, M., Escaff, D. & Clerc, M., 2014 Strong interaction between plants induces circular barren patches: fairy circles. *Philos. Trans. R. Soc. Lond. A* .
- [227] R.Martinez-Garcia, Calabrese, J., Hernandez-Garcia, E. & Lopez, C., 2014 Minimal mechanisms for vegetation patterns in semiarid regions. *Philos. Trans. R. Soc. Lond. A* .